

Immunization Timing

Age 2 months	
DTaP	
HepB*	
Hib	
PCV (Pneumo)	
Polio (IPV)	

Age 4 months	Interval from previous dose
DTaP	1-2 months
HepB*	1-2 months
Hib	1-2 months
PCV (Pneumo)	1-2 months
Polio (IPV)	1-2 months

Age 6 months	Interval from previous dose
DTaP	1-2 months
Hib [†]	1-2 months
PCV (Pneumo)	1-2 months
Polio (IPV)	2-14 months

Age 12 months	Interval from previous dose
HepB*	2-12 months and at least 4 months after 1st dose
Hib	2-8 months and on or after first birthday
PCV (Pneumo)	6-8 weeks and on or after first birthday
MMR	On or after first birthday
Varicella	On or after first birthday

Age 15 months	Interval from previous dose
DTaP	6-12 months

Age 2 years	Interval from previous dose
HepA	2nd dose 6-12 months after first

Before Kindergarten

IPV, DTaP, and MMR #2.
Also, unless already given,
HepB and Varicella.

Before 7th Grade

Unless already given,
HepB series, MMR #2, Varicella,
and Td booster.



CALIFORNIA KIDS
Love them. Immunize them.

This is a suggested schedule. For alternatives and details, consult the latest "Recommended Childhood Immunization Schedule, United States."

* HepB series can be started at birth, the second dose given at 1-2 months and the third dose at 6 months.

† This dose can be skipped if Pedvax Hib® and/or Comvax® (combined Hib and HepB vaccine) are used exclusively for the infant doses.